



Important Correction Notice

Demystifying Drug Dosing in Obese Patients by Brandon R. Shank and David E. Zimmerman

The publishers wish to inform you of a correction in Chapter 4, page 84.

Page 84, Chapter 4, Critical Care, under the heading “Ibutilide”

In the published book, dosing appears as “10 mg (one vial).” The corrected version of the paragraph is as follows:

Ibutilide is a Vaughan-Williams Class III antiarrhythmic indicated for the rapid conversion of atrial fibrillation or atrial flutter.³⁵ The dosing for ibutilide recommends **1 mg** IV infusion over 10 minutes for patients who weigh 60 kg or more.³⁵ A second infusion of **1 mg** can be given 10 minutes later if the arrhythmias did not terminate. No other data are available for dosing in obese patients, and the above dosing of **1 mg** should be used.

For print copy purchasers:

If you have a print copy of *Demystifying Drug Dosing in Obese Patients*, we ask that you return the copy to ASHP. ASHP will send you a corrected copy at no charge. For shipping the book to ASHP, please call 866-279-0681 or email us at publications@ashp.org for a prepaid shipping label.

For ebook purchasers:

Please delete your current ebook file for this title. To receive a corrected ebook file, go to your ebook library at ebooks.ashp.org.